arXiv:1507.03272v2 [math.CV] 26 Mar 2016

# EXPLICIT HODGE DECOMPOSITION ON RIEMANN SURFACES

GENNADI M. HENKIN AND PETER L. POLYAKOV

ABSTRACT. We present a construction of an explicit Hodge decomposition for $\bar{\partial}$-operator on Riemann surfaces.

## 1. INTRODUCTION

Classical Hodge decomposition on the space $Z^{(0,1)}(V) \subset \mathcal{E}^{(0,1)}(V)$ of smooth $\bar{\partial}$-closed $(0,1)$-forms on a smooth algebraic curve $V \subset \mathbb{CP}^n$, $(n \geq 2)$ with metric induced by Fubini-Study metric of $\mathbb{CP}^n$, has the following form

**Hodge Theorem .** *(*[Ho, Wy1, Wy2, Kd1]*) For any form $\phi \in Z^{(0,1)}(V)$ there exists a unique Hodge decomposition:*

$$\phi = \bar{\partial} R_1[\phi] + H_1[\phi], \tag{1.1}$$

*where $H_1$ is the orthogonal projection operator from $Z^{(0,1)}(V)$ onto the subspace $\mathcal{H}^{(0,1)}(V)$ of antiholomorphic $(0,1)$-forms on $V$, $R_1 = \bar{\partial}^* G_1$, $\bar{\partial}^* : \mathcal{E}^{(0,1)}(V) \to \mathcal{E}^{(0,0)}(V)$, where $\bar{\partial}^* = - * \bar{\partial} *$ is the Hodge dual operator for $\bar{\partial}$, $*$ is the Hodge operator, and $G_1$ is the Hodge-Green operator for Laplacian $\triangle = \bar{\partial}\bar{\partial}^* + \bar{\partial}^*\bar{\partial}$ on $V$.*

Hodge Theorem was proved by Hodge in [Ho] using the Fredholm's theory of integral equations. Weyl used his method of orthogonal projection from [Wy1] to correct and simplify the Hodge's proof in [Wy2], and was followed by Kodaira in [Kd1], who also used Weyl's method of orthogonal projection. However, the Hodge Theorem, as it is formulated above and in [Dl, BDIP], is not explicit enough for some applications. This disadvantage was pointed out by Griffiths and Harris in ([GH] §0.6), where the authors remarked that "the Hilbert space method has the disadvantage of not giving us the Green's operator" in the form of an integral operator with "a beautiful kernel on $M \times M$ with certain singularities along the diagonal".

A specific problem that we have in mind is an explicit solution of the inverse conductivity problem on a bordered Riemann surface, in which the conductivity function has to be reconstructed from the Dirichlet-to-Neumann map on its boundary (see [C], [HN]), in more general setting going back to [Ge]. We notice that article [HN] of Henkin and Novikov on this subject was motivated by article [HP1] by the authors of the present article. In [HP4] we made the first step toward explicit solution of the inverse conductivity problem by constructing an explicit Hodge-type decomposition for *$\bar{\partial}$-closed residual currents* of homogeneity zero on reduced complete intersections in $\mathbb{CP}^n$. In the present article using Theorem 1 from [HP4] we construct an explicit formula for operator $R_1$ in (1.1) assuming the knowledge of operator $H_1$. A problem that is definitely worth considering is the construction of an explicit form of $H_1$. Our choice of Riemann surfaces is motivated by the application mentioned above, though we consider the generalization of Theorem 1 from [HP4] to locally complete intersections as another interesting and important task.

---

2010 *Mathematics Subject Classification.* Primary: 14C30, 32S35, 32C30.

*Key words and phrases.* $\bar{\partial}$-operator, Riemann surface, Hodge decomposition.

Second author was partially supported by the NEUP program of the Department of Energy.

The main result of the present article is the construction in Theorem 2 of an explicit Hodge decomposition for $\bar{\partial}$-closed forms on an arbitrary Riemann surface assuming the knowledge of the Hodge projection. This construction is based on a generalized version of Theorem 1 from [HP4], which is presented in section 2 and covers the case of arbitrary homogeneity. The decompositions in Theorem 1 and in the propositions below are explicit in the sense that they depend only on the equations from (2.1) describing $V$ as a subvariety of $\mathbb{CP}^n$, and are defined by explicit integral operators with singular kernels of the Coleff-Herrera [CH] and of the Cauchy-Weil-Leray types [Wi, L].

Construction of integral formulas on $\mathbb{CP}^n$ with application to complex Radon transform was initiated in [HP1] and [Bn]. Application of such formulas to solution of $\bar{\partial}$-equation on singular analytic spaces was initiated in [HP2] and motivated further work in this direction (see for example [AS, FG, Sa]). The development of specific residual formulas in [HP4] and in Theorem 1 above has a long history going back to Poincare [P], Leray [L], Grothendieck [Gr], Herrera-Lieberman [HL], Dolbeault [Do], and Coleff and Herrera [CH]. As it is pointed out in [CH], the authors' work was conceived on the one hand as a generalization of the theory of residues of meromorphic forms and of the Grothendieck's theory of residues presented by Hartshorne [Ha1], and on the other hand of the work of Ramis and Ruget [RR] on dualizing complex in analytic geometry.

The modern development of the formulas of Cauchy-Weil-Leray type was initiated in [Kp], [Po], [Li], [O].

As a preliminary step in the construction of the explicit Hodge decomposition of Theorem 2 we use Theorem 1 and construct in Proposition 3.2 an intermediate explicit Hodge-type decomposition on an arbitrary compact Riemann surface $\mathcal{X}$, not necessarily embeddable into $\mathbb{CP}^2$. According to a classical result (see [Kd2]), going back to Gauss and Riemann, such surface admits an embedding as an algebraic submanifold in $\mathbb{CP}^3$. This embedding can be composed with a generic projection on $\mathbb{CP}^2$ to produce an immersion into $\mathbb{CP}^2$ with only nodes as singularities (see [GH, Ha2]). Then we use the Hodge-type decomposition of Theorem 1 on the image $\mathcal{C}$ of this immersion, which we lift and appropriately modify on $\mathcal{X}$. An important role in this construction is played by Proposition 3.1, in which we establish an isomorphism between the residual cohomologies on $\mathcal{C}$ and the cohomologies of the structural sheaf of $\mathcal{C}$. This proposition might be considered as a step in constructing a Hodge-type decomposition of cohomologies on curves with singularities, following direction of [Gr],[Ha2], and [Dl]. We notice that the proof of Theorem 2 generalizes verbatim to the case of a nonsingular projective complete intersection.

In section 4 using Theorems 1 and 2 we obtain explicit formulas for solutions of $\bar{\partial}$-equation and present two explicit versions of the *Hodge-Kodaira Vanishing Theorem* for open Riemann surfaces.

## 2. GENERALIZED VERSION OF THEOREM 1 FROM [HP4].

Below we present a generalized version of Theorem 1 from [HP4], which gives a Hodge-type decomposition of residual currents of arbitrary homogeneity on reduced complete intersections in $\mathbb{CP}^n$. Before formulating this version we introduce definitions from [HP3] and [HP4].

Let $V$ be a complete intersection subvariety

$$V = \{z \in \mathbb{CP}^n :\ P_1(z) = \cdots = P_m(z) = 0\} \tag{2.1}$$

of dimension $n-m$ in $\mathbb{CP}^n$ defined by a collection $\{P_k\}_{k=1}^m$ of homogeneous polynomials. Let

$$\{U_\alpha = \{z \in \mathbb{CP}^n :\ z_\alpha \neq 0\}\}_{\alpha=0}^n$$

be the standard covering of $\mathbb{CP}^n$, and let

$$\mathbf{F}^{(\alpha)}(z) = \begin{bmatrix} F_1^{(\alpha)}(z) \\ \vdots \\ F_m^{(\alpha)}(z) \end{bmatrix} = \begin{bmatrix} P_1(z)/z_\alpha^{\deg P_1} \\ \vdots \\ P_m(z)/z_\alpha^{\deg P_m} \end{bmatrix}$$

be collections of nonhomogeneous polynomials satisfying

$$\mathbf{F}^{(\alpha)}(z) = A_{\alpha\beta}(z)\cdot\mathbf{F}^{(\beta)}(z) = \begin{bmatrix} (z_\beta/z_\alpha)^{\deg P_1} & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & (z_\beta/z_\alpha)^{\deg P_m} \end{bmatrix}\cdot\mathbf{F}^{(\beta)}(z)$$

on $U_{\alpha\beta} = U_\alpha \cap U_\beta$.

Following [Gr] and [Ha2] we consider a line bundle $\mathcal{L}$ on $V$ with transition functions

$$l_{\alpha\beta}(z) = \det A_{\alpha\beta} = \left(\frac{z_\beta}{z_\alpha}\right)^{\sum_{k=1}^m \deg P_k}$$

on $U_{\alpha\beta}$ and the *dualizing bundle* on the complete intersection subvariety $V$

$$\omega_V^\circ = \omega_{\mathbb{CP}^n}\otimes\mathcal{L}, \tag{2.2}$$

where $\omega_{\mathbb{CP}^n}$ is the canonical bundle on $\mathbb{CP}^n$.

For $q = 1,\dots,n-m$ we denote by $\mathcal{E}^{(n,n-m-q)}(V,\mathcal{L}(-\ell)) = \mathcal{E}^{(0,n-m-q)}(V,\omega_V^\circ(-\ell))$ the space of $C^\infty$ differential forms of bidegree $(n,n-m-q)$ with coefficients in $\mathcal{L}\otimes\mathcal{O}(-\ell)$, i.e. the space of collections of forms

$$\left\{\gamma_\alpha \in \mathcal{E}^{(n,n-m-q)}(U_\alpha)\right\}_{\alpha=0}^n$$

satisfying

$$\gamma_\alpha = l_{\alpha\beta}\cdot\left(\frac{z_\beta}{z_\alpha}\right)^{-\ell}\gamma_\beta + \sum_{k=1}^m F_k^{(\alpha)}\cdot\gamma_k^{\alpha\beta} \text{ on } U_\alpha\cap U_\beta. \tag{2.3}$$

Then, following [CH, HP3, Pa] we define residual currents and $\bar\partial$-closed residual currents on $V$. By a residual current $\phi \in C_R^{(0,q)}(V,\mathcal{O}(\ell))$ of homogeneity $\ell$ we call a collection $\left\{\Phi_\alpha^{(0,q)}\right\}_{\alpha=0}^n$ of $C^\infty$ differential forms satisfying equalities

$$\Phi_\alpha = \left(\frac{z_\beta}{z_\alpha}\right)^\ell\Phi_\beta + \sum_{k=1}^m F_k^{(\alpha)}\cdot\Omega_k^{(\alpha\beta)} \text{ on } U_\alpha\cap U_\beta, \tag{2.4}$$

acting on $\gamma \in \mathcal{E}^{(n,n-m-q)}(V,\mathcal{L}(-\ell))$ by the formula

$$\langle\phi,\gamma\rangle = \sum_\alpha\int_{U_\alpha}\vartheta_\alpha\gamma_\alpha\wedge\Phi_\alpha\bigwedge_{k=1}^m\bar\partial\frac{1}{F_k^{(\alpha)}} \overset{\text{def}}{=} \lim_{t\to0}\sum_\alpha\int_{T_\alpha^{\epsilon(t)}}\vartheta_\alpha\frac{\gamma_\alpha\wedge\Phi_\alpha}{\prod_{k=1}^m F_k^{(\alpha)}}, \tag{2.5}$$

where $\{\vartheta_\alpha\}_{\alpha=0}^n$ is a partition of unity subordinate to the covering $\{U_\alpha\}_{\alpha=0}^n$, and the limit in the right-hand side of (2.5) is taken along an admissible path in the sense of Coleff-Herrera [CH], i.e. an analytic map $\epsilon : [0,1] \to \mathbb{R}^m$ satisfying conditions

$$\begin{cases} \lim_{t\to0}\epsilon_m(t) = 0, \\ \lim\limits_{t\to0}\dfrac{\epsilon_j(t)}{\epsilon_{j+1}^l(t)} = 0, \text{ for any } l\in\mathbb{N} \text{ and } j = 1,\dots,m-1, \end{cases} \tag{2.6}$$

and

$$T_\alpha^{\epsilon(t)} = \left\{z\in U_\alpha : \left|F_k^{(\alpha)}(z)\right| = \epsilon_k(t) \text{ for } k = 1,\dots,m\right\}. \tag{2.7}$$

A residual current $\phi$ of homogeneity $\ell$ we call $\bar\partial$-closed $\left(\text{denoted } \phi \in Z_R^{(0,q)}(V,\mathcal{O}(\ell))\right)$, if it satisfies the following condition

$$\bar\partial\Phi_\alpha = \sum_{k=1}^m F_k^{(\alpha)}\cdot\Omega_k^{(\alpha)} \text{ on } U_\alpha. \tag{2.8}$$

Below we present an extended version of Theorem 1 from [HP4] that is used in this article.

**Theorem 1.** *Let* $V \subset \mathbb{CP}^n$ *be a reduced complete intersection subvariety as in* (2.1). *Then*

(i) *for an arbitrary* $\phi \in Z_R^{(0,q)}\left(V, \mathcal{O}(\ell)\right)$ *the following representation holds*

$$\phi = \bar{\partial} I_q[\phi] + L_q[\phi], \tag{2.9}$$

*where* $L_q[\phi] = 0$ *if* $1 \leq q < n-m$, $L_{n-m}[\phi] \in Z_R^{(0,n-m)}\left(V, \mathcal{O}(\ell)\right)$ *is defined by formula*

$$L_{n-m}[\phi] = \sum_{0 \leq r \leq d-n-1-\ell} C(n,m,d,r) \lim_{t \to 0} \int_{\left\{|\zeta|=1, \{|P_k(\zeta)|=\epsilon_k(t)\}_{k=1}^m\right\}} \langle \bar{z} \cdot \zeta \rangle^r \cdot \frac{\phi(\zeta)}{\prod_{k=1}^m P_k(\zeta)}$$
$$\bigwedge \det \left[ \bar{z} \ \overbrace{Q(\zeta,z)}^{m} \ \overbrace{d\bar{z}}^{n-m} \right] \wedge \omega(\zeta) \tag{2.10}$$

*with* $d = \sum_{k=1}^m \deg P_k$, *and the current* $I_q[\phi] \in C^{(0,q-1)}\left(V, \mathcal{O}(\ell)\right)$ *is defined by formula*

$$I_q[\phi] = C(n,q,m) \lim_{t \to 0} \int_{\left\{|\zeta|=1, \{|P_k(\zeta)|=\epsilon_k(t)\}_{k=1}^m\right\}} \frac{\phi(\zeta)}{\prod_{k=1}^m P_k(\zeta)}$$
$$\bigwedge \det \left[ \frac{\bar{z}}{B^*(\zeta,z)} \ \frac{\bar{\zeta}}{B(\zeta,z)} \ \overbrace{Q(\zeta,z)}^{m} \ \overbrace{\frac{d\bar{z}}{B^*(\zeta,z)}}^{q-1} \ \overbrace{\frac{d\bar{\zeta}}{B(\zeta,z)}}^{n-m-q} \right] \wedge \omega(\zeta), \tag{2.11}$$

*where functions* $\left\{Q_k^i(\zeta,z)\right\}$ *for* $k = 1, \dots, m, \ i = 0, \dots, n$ *satisfy*

$$\begin{cases} P_k(\zeta) - P_k(z) = \sum_{i=0}^n Q_k^i(\zeta,z) \cdot (\zeta_i - z_i), \\ Q_k^i(\lambda\zeta, \lambda z) = \lambda^{\deg P_k - 1} \cdot Q_k^i(\zeta,z) \text{ for } \lambda \in \mathbb{C}, \end{cases} \tag{2.12}$$

*and*

$$B^*(\zeta,z) = \sum_{j=0}^n \bar{z}_j \cdot (\zeta_j - z_j), \qquad B(\zeta,z) = \sum_{j=0}^n \bar{\zeta}_j \cdot (\zeta_j - z_j);$$

(ii) *a* $\bar{\partial}$*-closed residual current* $\phi \in Z_R^{(0,n-m)}\left(V, \mathcal{O}(\ell)\right)$ *is* $\bar{\partial}$*-exact, i.e. there exists a current* $\psi \in C^{(0,n-m-1)}\left(V, \mathcal{O}(\ell)\right)$ *such that* $\phi = \bar{\partial}\psi$, *iff*

$$L_{n-m}[\phi] = 0. \tag{2.13}$$

We present here a sketch of proof of the Theorem. In this sketch we are concerned with extending the validity of all lemmas and propositions comprising the proof of Theorem 1 in [HP4] to the case of nonzero homogeneity.

In the lemma below we prove that the operators $L$ and $I$ defined in [HP4] preserve homogeneity, validating equalities (2.6), (2.11) and Proposition 2.3 from [HP4] in the case of nonzero homogeneity.

**Lemma 2.1.** *If* $\phi \in C_R^{(0,q)}(V, \mathcal{O}(\ell))$ *is a residual current defined by a differential form* $\Phi$ *in a neighborhood of* $V$ *satisfying*

$$\Phi(\lambda\zeta) = \lambda^{\ell} \cdot \Phi(\zeta), \tag{2.14}$$

*then for the forms* $L_{n-m}\left[\Phi\right]$ *and* $I_q\left[\Phi\right]$ *defined by operators* $L_{n-m}$ *and* $I_q$ *in formulas (3.24) and (4.22) of* [HP4] *respectively, the following equalities hold*

$$\begin{aligned} L_{n-m}\left[\Phi\right](\lambda z) &= \lambda^{\ell} \cdot L_{n-m}\left[\Phi\right](z), \\ I_q\left[\Phi\right](\lambda z) &= \lambda^{\ell} \cdot I_q\left[\Phi\right](z). \end{aligned} \tag{2.15}$$

*Proof.* To prove the preservation of homogeneity for operator $L_{n-m}$ we use the following equality for $|\lambda| = 1$, after changing variables to $\zeta = \lambda w$

$$
\begin{aligned}
&\int_{\{|\zeta|=1,\{|P_k(\zeta)|=\epsilon_k(t)\}_{k=1}^m\}} \langle \bar\lambda \bar z \cdot \zeta\rangle^r \cdot \frac{\Phi(\zeta)}{\prod_{k=1}^m P_k(\zeta)} \bigwedge \det \left[ \bar\lambda \bar z \; \overbrace{Q(\zeta, \lambda z)}^{m} \; \overbrace{\bar\lambda d\bar z}^{n-m} \right] \wedge \omega(\zeta) \\
&= \int_{\{|\zeta|=1,\{|P_k(w)|=\epsilon_k(t)\}_{k=1}^m\}} \langle \bar\lambda \bar z \cdot \lambda w\rangle^r \cdot \frac{\Phi(\lambda w)}{\prod_{k=1}^m P_k(\lambda w)} \bigwedge \det \left[ \bar\lambda \bar z \; \overbrace{Q(\lambda w, \lambda z)}^{m} \; \overbrace{\bar\lambda d\bar z}^{n-m} \right] \wedge \omega(\lambda w) \\
&\qquad = \lambda^{\ell-d}\bar\lambda^{n-m+1}\lambda^{d-m}\lambda^{n+1} \int_{\{|\zeta|=1,\{|P_k(w)|=\epsilon_k(t)\}_{k=1}^m\}} \langle \bar z \cdot w\rangle^r \cdot \frac{\Phi(w)}{\prod_{k=1}^m P_k(w)} \\
&\qquad\qquad \bigwedge \det \left[ \bar\lambda \bar z \; \overbrace{Q(w, z)}^{m} \; \overbrace{d\bar z}^{n-m} \right] \wedge \omega(w) \\
&= \lambda^{\ell} \int_{\{|\zeta|=1,\{|P_k(w)|=\epsilon_k(t)\}_{k=1}^m\}} \langle \bar z \cdot w\rangle^r \cdot \frac{\Phi(w)}{\prod_{k=1}^m P_k(w)} \bigwedge \det \left[ \bar\lambda \bar z \; \overbrace{Q(w, z)}^{m} \; \overbrace{d\bar z}^{n-m} \right] \wedge \omega(w),
\end{aligned}
$$

where $d = \sum_{k=1}^m \deg P_k$.

Similarly, we obtain equality

$$
\begin{aligned}
&\int_{\{|\zeta|=1,\{|P_k(w)|=\epsilon_k(t)\}_{k=1}^m\}} \frac{\Phi(\lambda w)}{\prod_{k=1}^m P_k(\lambda w)} \\
&\quad \bigwedge \det \left[ \frac{\bar\lambda \bar z}{B^*(\lambda w, \lambda z)} \; \frac{\bar\lambda \bar w}{B(\lambda w, \lambda z)} \; \overbrace{Q(\lambda w, \lambda z)}^{m} \; \overbrace{\frac{\bar\lambda d\bar z}{B^*(\lambda w, \lambda z)}}^{q-1} \; \overbrace{\frac{\bar\lambda d\bar w}{B(\lambda w, \lambda z)}}^{n-m-q} \right] \wedge \omega(\lambda w) \\
&\qquad = \lambda^{\ell-d}\lambda^{d-m}\bar\lambda^{n-m+1}\lambda^{n+1} \int_{\{|\zeta|=1,\{|P_k(w)|=\epsilon_k(t)\}_{k=1}^m\}} \frac{\Phi(w)}{\prod_{k=1}^m P_k(w)} \\
&\qquad\quad \bigwedge \det \left[ \frac{\bar z}{B^*(w, z)} \; \frac{\bar w}{B(w, z)} \; \overbrace{Q(w, z)}^{m} \; \overbrace{\frac{d\bar z}{B^*(w, z)}}^{q-1} \; \overbrace{\frac{d\bar w}{B(w, z)}}^{n-m-q} \right] \wedge \omega(w) \\
&\qquad\qquad = \lambda^{\ell} \int_{\{|\zeta|=1,\{|P_k(w)|=\epsilon_k(t)\}_{k=1}^m\}} \frac{\Phi(w)}{\prod_{k=1}^m P_k(w)} \\
&\qquad\qquad\quad \bigwedge \det \left[ \frac{\bar z}{B^*(w, z)} \; \frac{\bar w}{B(w, z)} \; \overbrace{Q(w, z)}^{m} \; \overbrace{\frac{d\bar z}{B^*(w, z)}}^{q-1} \; \overbrace{\frac{d\bar w}{B(w, z)}}^{n-m-q} \right] \wedge \omega(w).
\end{aligned}
$$

□

To include the case of nonzero homogeneity Lemmas 3.2 and 3.3 in [HP4] have to be reformulated. In particular, Lemma 3.2 has to be replaced by the following Lemma.

**Lemma 2.2.** *Let $V\subset\mathbb{CP}^n$ be a reduced complete intersection subvariety as in* (2.1)*, let $U\supset V$ be an open neighborhood of $V$ in $\mathbb{CP}^n$, and let $\Phi\in\mathcal{E}_c^{(0,n-m)}(U\cap U_\alpha)$ be a differential form of homogeneity $\ell$ on $U\cap U_\alpha$ for some $\alpha\in(0,\dots,n)$.*

*Then formula*

$$\lim_{t\to 0}\int_{\left\{|\zeta|=\tau,\{|P_k(\zeta)|=\epsilon_k(t)\}_{k=1}^m\right\}}\langle\bar z\cdot\zeta\rangle^r\cdot\frac{\Phi(\zeta)}{\prod_{k=1}^m P_k(\zeta)}\wedge\det\left[\bar z\ \overbrace{Q(\zeta,z)}^{m}\ \overbrace{d\bar z}^{n-m}\right]\wedge\omega(\zeta),\tag{2.16}$$

*where $\{\epsilon_k(t)\}_{k=1}^m$ is an admissible path, defines a differential form of homogeneity $\ell$ on $U$, real analytic with respect to $z$.*

*If $\Phi(\zeta)=\sum_{k=1}^m F_k^{(\alpha)}(\zeta)\Omega_k(\zeta)$ with $\Omega_k\in\mathcal{E}_c^{(0,n-m)}(U\cap U_\alpha)$, then the limit in* (2.16) *is equal to zero.*

*Proof.* Preservation of homogeneity follows from the first equality in (2.15) in Lemma 2.1. As in Lemma 3.2 in [HP4] without loss of generality we can consider only the case $\alpha=0$, i.e. $\Phi\in\mathcal{E}_c^{(0,n-m)}(U\cap\{\zeta_0\neq 0\})$. Then formula (3.15) in [HP4] for a form $\Phi$ satisfying equality (2.14) has to be replaced by the following formula

$$\begin{aligned}
&\int_{\left\{|\zeta|=\tau,\{|P_k(\zeta)|=\epsilon_k(t)\}_{k=1}^m\right\}}\left(\bar z_0+\sum_{j=1}^n\bar z_j\cdot w_j\right)^r\cdot\frac{\Phi(\zeta)}{\prod_{k=1}^m P_k(\zeta)}\\
&\qquad\wedge\det\left[\bar z\ \overbrace{Q(\zeta,z)}^{m}\ \overbrace{d\bar z}^{n-m}\right]\wedge\left(\zeta_0^{n+r}d\zeta_0\right)\bigwedge_{j=1}^n dw_j\\
&=\int_{\left\{|\zeta|=\tau,\{|P_k(\zeta)|=\epsilon_k(t)\}_{k=1}^m\right\}}\left(\bar z_0+\sum_{j=1}^n\bar z_j\cdot w_j\right)^r\cdot\frac{\Phi(\zeta)}{\prod_{k=1}^m P_k(\zeta)}\\
&\qquad\wedge\det\left[\bar z\ \overbrace{Q(\zeta,z)}^{m}\ \overbrace{d\bar z}^{n-m}\right]\wedge\left(\zeta_0^{n+r}d\zeta_0\right)\bigwedge_{j=1}^n dw_j\\
&=\int_{\left\{|\zeta|=\tau,\{|F_k(w)|\cdot\chi_k(w)=\epsilon_k(t)\}_{k=1}^m\right\}}\left(\bar z_0+\sum_{j=1}^n\bar z_j\cdot w_j\right)^r\left(\zeta_0^{n+r+\ell-\sum_{k=1}^m\deg P_k}d\zeta_0\right)\\
&\qquad\times\frac{\Phi(w)}{\prod_{k=1}^m F_k(w)}\wedge\det\left[\bar z\ \overbrace{Q(e^{i\phi_0},w,z)}^{m}\ \overbrace{d\bar z}^{n-m}\right]\bigwedge_{j=1}^n dw_j\\
&=i\int_0^{2\pi}e^{i\left(n+r+\ell-\sum_{k=1}^m\deg P_k+1\right)\phi_0}d\phi_0\int_{\left\{w\in U_0,\{|F_k(w)|\cdot\chi_k(w)=\epsilon_k(t)\}_{k=1}^m\right\}}\left(\bar z_0+\sum_{j=1}^n\bar z_j\cdot w_j\right)^r\\
&\qquad\times\rho_0(w)^{n+r+\ell-\sum_{k=1}^m\deg P_k+1}\cdot\frac{\Phi(w)}{\prod_{k=1}^m F_k(w)}\wedge\det\left[\bar z\ \overbrace{Q(e^{i\phi_0},w,z)}^{m}\ \overbrace{d\bar z}^{n-m}\right]\bigwedge_{j=1}^n dw_j,
\end{aligned}\tag{2.17}$$

where we used nonhomogeneous coordinates $\{w_i=\zeta_i/\zeta_0\}_{i=1}^n$ in the subset $\mathbb{S}^{2n+1}(\tau)\setminus\{\zeta_0=0\}$ of the sphere $\mathbb{S}^{2n+1}(\tau)$ of radius $\tau$ in $\mathbb{C}^{n+1}$, notation $\zeta_0=\rho_0(w)\cdot e^{i\phi_0}$ with

$$\rho_0(w)=\frac{\tau}{\sqrt{1+\sum_{i=1}^n|w_i|^2}}$$

on $\mathbb{S}^{2n+1}(\tau)$, nonhomogeneous polynomials

$$F_k(w) = P_k(\zeta)/\zeta_0^{\deg P_k}$$

in $\mathbb{S}^{2n+1}(\tau)\setminus\{\zeta_0 = 0\}$, and equality (2.14).

The last statement of the Lemma follows as in Lemma 3.2 of [HP4] from application of Theorem 1.7.6(2) in [CH] to the interior integral in the right-hand side of (2.17). □

Lemma 3.3 has to be replaced by the following Lemma.

**Lemma 2.3.** *Let $\phi \in Z_R^{(0,q)}(V, \mathcal{O}(\ell))$ be a $\bar\partial$-closed residual current defined by a collection of forms $\left\{\Phi_\alpha^{(0,n-m)}\right\}_{\alpha=0}^{n}$ of homogeneity $\ell$ on a neighborhood $U$ of the reduced subvariety $V$ as in* (2.1) *satisfying* (2.4) *and* (2.8)*, and let $\Phi(\zeta) = \sum_{\alpha=0}^{n} \vartheta_\alpha(\zeta)\Phi_\alpha(\zeta)$ be a differential form of homogeneity $\ell$ on $U$.*

*Then for an arbitrary $\gamma \in \mathcal{E}^{(n,0)}\left(V, \mathcal{L}(-\ell)\right)$ the equality*

$$\lim_{\tau\to 0}\int_{T_\beta^{\delta(\tau)}} \frac{\gamma(z)}{\prod_{k=1}^{m} F_k^{(\beta)}(z)} \bigwedge \left(\lim_{t\to 0}\int_{\left\{|\zeta|=1,\{|P_k(\zeta)|=\epsilon_k(t)\}_{k=1}^{m}\right\}} \langle \bar z\cdot\zeta\rangle^r \times \frac{\Phi(\zeta)}{\prod_{k=1}^{m} P_k(\zeta)} \wedge \det\left[\bar z\ \overbrace{Q(\zeta,z)}^{m}\ \overbrace{d\bar z}^{n-m}\right] \wedge \omega(\zeta)\right) = 0 \tag{2.18}$$

*holds unless*

$$r \le \sum_{k=1}^{m} \deg P_k - \ell - n - 1. \tag{2.19}$$

*Proof.* We notice that for all values of $a > 0$ and $\epsilon = (\epsilon_1, \dots, \epsilon_k)$ the sets

$$S(a) = \left\{\zeta \in \mathbb{S}^{2n+1}(a) : \left\{|P_k(\zeta)| = \epsilon_k \cdot a^{\deg P_k}\right\}_{k=1}^{m}\right\}$$

are real analytic subvarieties of $\mathbb{S}^{2n+1}(a)$ of real dimension $2n+1-m$ satisfying

$$c \cdot a^{2n+1-m} \cdot \text{Volume}\left(S(1)\right) < \text{Volume}_{2n+1-m}\left(S(a)\right) < C \cdot a^{2n+1-m} \cdot \text{Volume}\left(S(1)\right). \tag{2.20}$$

We denote

$$\Phi_\alpha(\zeta, z) = \langle \bar z\cdot\zeta\rangle^r \cdot \frac{\Phi_\alpha(\zeta)}{\prod_{k=1}^{m} P_k(\zeta)} \det\left[\bar z\ \overbrace{Q(\zeta,z)}^{m}\ \overbrace{d\bar z}^{n-m}\right] \wedge \omega(\zeta),$$

and apply the Stokes' formula to the differential form

$$\beta(\zeta, z) = \sum_{\alpha=0}^{n} \vartheta_\alpha(\zeta)\Phi_\alpha(\zeta, z) = \sum_{\alpha=0}^{n} \langle \bar z\cdot\zeta\rangle^r \cdot \frac{\vartheta_\alpha(\zeta)\Phi_\alpha(\zeta)}{\prod_{k=1}^{m} P_k(\zeta)} \det\left[\bar z\ \overbrace{Q(\zeta,z)}^{m}\ \overbrace{d\bar z}^{n-m}\right] \wedge \omega(\zeta) \tag{2.21}$$

on the variety

$$\left\{\zeta \in \mathbb{C}^{n+1} : \left\{|P_k(\zeta)| = \epsilon_k \cdot |\zeta|^{\deg P_k}\right\}_{k=1}^{m},\ a < |\zeta| < 1\right\}$$

with the boundary

$$\left\{\zeta : |\zeta| = a,\ \left\{|P_k(\zeta)| = \epsilon_k \cdot a^{\deg P_k}\right\}_{k=1}^{m}\right\} \bigcup \left\{\zeta : |\zeta| = 1,\ \{|P_k(\zeta)| = \epsilon_k\}_{k=1}^{m}\right\}.$$

Then using equality (2.8) we obtain the equality

$$\begin{aligned}\int_{\left\{|\zeta|=1,\{|P_k(\zeta)|=\epsilon_k(t)\}_{k=1}^m\right\}}\beta(\zeta,z)-\int_{\left\{|\zeta|=a,\ \left\{|P_k(\zeta)|=\epsilon_k(t)\cdot a^{\deg P_k}\right\}_{k=1}^m\right\}}\beta(\zeta,z)\\ =\sum_{\alpha=0}^{n}\sum_{j=1}^{m}\int_a^1 d\tau\int_{\left\{|\zeta|=\tau,\left\{|P_k(\zeta)|=\epsilon_k(t)\cdot\tau^{\deg P_k}\right\}_{k=1}^m\right\}}F_j^{(\alpha)}(\zeta)\cdot\beta_j^{(\alpha)}(\zeta,z)\\ +\sum_{\alpha=0}^{n}\int_a^1 d\tau\int_{\left\{|\zeta|=\tau,\left\{|P_k(\zeta)|=\epsilon_k(t)\cdot\tau^{\deg P_k}\right\}_{k=1}^m\right\}}d|\zeta|\lrcorner\left(\bar\partial\vartheta_\alpha(\zeta)\wedge\Phi_\alpha(\zeta,z)\right)\end{aligned}\tag{2.22}$$

for arbitrary $t$ and $0<a<1$.

Using estimate (2.20) and the homogeneity property

$$\Phi_\beta(t\cdot\zeta)=\bar t^{-(n-m)}\cdot t^{\ell}\cdot\Phi_\beta(\zeta)\tag{2.23}$$

of the coefficients of $\Phi^{(0,n-m)}$ from Proposition 1.1 in [HP1] we obtain that if

$$r+n+1+\ell-\sum_{k=1}^m\deg P_k>0,\tag{2.24}$$

then

$$\left|\int_{\left\{|\zeta|=a,\ \left\{|P_k(\zeta)|=\epsilon_k(t)\cdot a^{\deg P_k}\right\}_{k=1}^m\right\}}\beta(\zeta,z)\right|<C'(\epsilon)\cdot a^{r+2n+1-m-(n-m)+\ell-\sum_{k=1}^m\deg P_k}\longrightarrow 0$$

as $t$ is fixed and $a\to 0$.

For the first sum of integrals in the right-hand side of (2.22) we have

$$\begin{aligned}\left|\int_a^1 d\tau\int_{\left\{|\zeta|=\tau,\ \left\{|P_k(\zeta)|=\epsilon_k(t)\cdot\tau^{\deg P_k}\right\}_{k=1}^m\right\}}F_j^{(\alpha)}(\zeta)\cdot\beta_j^{(\alpha)}(\zeta,z)\right|\\ <C\int_a^1 d\tau\cdot\tau^{r+2n+1-m-(n-m)+\ell-\sum_{k=1}^m\deg P_k}\\ \times\int_{\left\{|\zeta|=\tau,\ \left\{|P_k(\zeta)|=\epsilon_k(t)\cdot\tau^{\deg P_k}\right\}_{k=1}^m\right\}}F_j^{(\alpha)}(\zeta)\cdot\left(d|\zeta|\lrcorner\beta_j^{(\alpha)}(\zeta,z)\right)\\ <C'(\epsilon)\frac{\left(1-a^{r+n+2+\ell-\sum_{k=1}^m\deg P_k}\right)}{r+n+2+\ell-\sum_{k=1}^m\deg P_k}\longrightarrow 0\end{aligned}\tag{2.25}$$

as $t \to 0$, since $a < 1$, condition (2.24) is satisfied, and $C'(\epsilon) \to 0$ as $t \to 0$ by Lemma 2.2.

For the second sum of integrals in the right-hand side of (2.22) we use equality

$$
\begin{aligned}
&\lim_{t\to0} \sum_{\left\{\beta:\, U_\beta\cap U_\alpha\neq\emptyset\right\}} \int_a^1 d\tau \int_{\left\{\zeta_\alpha\neq0,|\zeta|=\tau,\left\{|P_k(\zeta)|=\epsilon_k(t)\cdot\tau^{\deg P_k}\right\}_{k=1}^m\right\}} d|\zeta|\rfloor\left(\bar\partial\vartheta_\beta(\zeta)\wedge\Phi_\beta(\zeta,z)\right)\\
&= \lim_{t\to0}\int_a^1 d\tau \sum_{\left\{\beta:\, U_\beta\cap U_\alpha\neq\emptyset\right\}} \int_{\left\{\zeta_\alpha\neq0,|\zeta|=\tau,\left\{|P_k(\zeta)|=\epsilon_k(t)\cdot\tau^{\deg P_k}\right\}_{k=1}^m\right\}} \langle\bar z\cdot\zeta\rangle^r\\
&\times d|\zeta|\rfloor\left(\bar\partial\vartheta_\beta(\zeta)\wedge\frac{\Phi_\beta(\zeta)}{\left[\prod_{k=1}^m P_k(\zeta)\right]\big|_\beta}\bigwedge\det\left[\bar z\ \overbrace{Q(\zeta,z)}^{m}\ \overbrace{d\bar z}^{n-m}\right]\wedge\omega(\zeta)\right)\\
&= \lim_{t\to0}\int_a^1 d\tau \sum_{\left\{\beta:\, U_\beta\cap U_\alpha\neq\emptyset\right\}} \int_{\left\{\zeta_\alpha\neq0,|\zeta|=\tau,\left\{|P_k(\zeta)|=\epsilon_k(t)\cdot\tau^{\deg P_k}\right\}_{k=1}^m\right\}} \langle\bar z\cdot\zeta\rangle^r\\
&\times d|\zeta|\rfloor\left(\bar\partial\vartheta_\beta(\zeta)\big|_{U_\alpha}\wedge\frac{\Phi_\alpha(\zeta)}{\left[\prod_{k=1}^m P_k(\zeta)\right]\big|_\alpha}\wedge\det\left[\bar z\ \overbrace{Q(\zeta,z)}^{m}\ \overbrace{d\bar z}^{n-m}\right]\wedge\omega(\zeta)\right)\\
&= \lim_{t\to0}\int_a^1 d\tau \int_{\left\{\zeta_\alpha\neq0,|\zeta|=\tau,\left\{|P_k(\zeta)|=\epsilon_k(t)\cdot\tau^{\deg P_k}\right\}_{k=1}^m\right\}} \langle\bar z\cdot\zeta\rangle^r\\
&\times d|\zeta|\rfloor\left(\left[\sum_{\left\{\beta:\, U_\beta\cap U_\alpha\neq\emptyset\right\}}\bar\partial\vartheta_\beta(\zeta)\big|_{U_\alpha}\right]\wedge\frac{\Phi_\alpha(\zeta)}{\left[\prod_{k=1}^m P_k(\zeta)\right]\big|_\alpha}\bigwedge\det\left[\bar z\ \overbrace{Q(\zeta,z)}^{m}\ \overbrace{d\bar z}^{n-m}\right]\wedge\omega(\zeta)\right)=0.
\end{aligned}
$$

which follows from equality (2.4), Lemma 2.2, and the transformation formula for the Grothendieck's residue from Proposition 4.2 in [HP1] (for isolated singularities see [GH, T]). □

This completes the sketch of proof of Theorem 1.

Below we formulate a corollary of Theorem 1, which will be used in what follows. This corollary gives an explicit form of the *Hodge-Kodaira Vanishing Theorem* for projective complete intersections (see III.7.15 in [Ha2] and §2.1 of [GH].

**Corollary 2.4.** *If homogeneity $\ell$ satisfies inequality*

$$\ell > d - n - 1, \tag{2.26}$$

*then the operator in the right-hand side of equality 2.10 is zero,*

$$\dim H_R^{n-m}\left(V,\mathcal{O}(\ell)\right) = 0, \tag{2.27}$$

*and for any $\phi \in Z_R^{(0,n-m)}\left(V,\mathcal{O}(\ell)\right)$*

$$\phi = \bar\partial I_{n-m}[\phi]. \tag{2.28}$$

## 3. EXPLICIT HODGE DECOMPOSITION ON NONSINGULAR ALGEBRAIC CURVES.

In order to construct an explicit Hodge decomposition on an arbitrary nonsingular algebraic curve $\mathcal{X}$ we use the existence of an immersion (see [Kd2], 1.4 in [GH], IV.3 in [Ha2])

$$\mathcal{X} \xrightarrow{\varrho} \mathbb{CP}^2, \tag{3.1}$$

such that $\mathcal{C} = \varrho(\mathcal{X})$ is a plane curve with at most nodes as singularities. Our construction of the sought decomposition on $\mathcal{X}$ will be based on a similar decomposition on $\mathcal{C}$, which exists according to

Theorem 1.

We construct a linear map $\varrho_* : \mathcal{E}^{(0,1)}(\mathcal{X}) \to Z^{(0,1)}_R(\mathcal{C})$. Here and below we use notations $\varrho_*$ and $\varrho^*$ for induced direct and respectively inverse maps on functions and differential forms. Since $\varrho$ is biholomorphic everywhere outside of nodal points, we have to describe $\varrho_*$ only in the neighborhoods of those points. Let $p \in \mathcal{C}$ be such nodal point with $p_1, p_2 \in \mathcal{X}$ such that $\varrho(p_1) = \varrho(p_2) = p$. Let $z_1, z_2$ be local coordinates at $p \in U \subset \mathbb{CP}^2$, such that $z_1(p) = z_2(p) = 0$, and such that

$$\mathcal{C} \cap U = \mathcal{C}_1 \cup \mathcal{C}_2,$$

where $\mathcal{C}_1 = \{q \in U : z_1(q) = 0\}$, and $\mathcal{C}_2 = \{q \in U : z_2(q) = 0\}$.

Let $\phi \in \mathcal{E}^{(0,1)}(\mathcal{X})$ be a smooth $\bar\partial$-closed form on $\mathcal{X}$ with local representations

$$\phi\Big|_{p_1} = \phi_1(z)d\bar z_1, \quad \phi\Big|_{p_2} = \phi_2(z)d\bar z_2$$

on $\mathcal{X}$. Let $\widetilde{\phi}_1, \widetilde{\phi}_2$ be extensions of $\phi_1$ and $\phi_2$ to $U$ such that

$$\widetilde{\phi}_j\Big|_{\mathcal{C}_j} = \phi_j, \quad \bar\partial_{z_k}\widetilde{\phi}_j = 0 \text{ for } k \neq j.$$

Then the differential form

$$\phi_* = \widetilde{\phi}_1 d\bar z_1 + \widetilde{\phi}_2 d\bar z_2 \tag{3.2}$$

defines a $\bar\partial$-closed residual current on $\mathcal{C}$ in the neighborhood $U \ni p$. Current $\phi_*$ is $\bar\partial$-exact in $U$ since for the functions $\widetilde{\psi}_1$ and $\widetilde{\psi}_2$ chosen so that

$$\bar\partial_{z_j}\widetilde{\psi}_j = \widetilde{\phi}_j, \quad \bar\partial_{z_k}\widetilde{\psi}_j = 0 \text{ for } k \neq j,$$

we will have

$$\bar\partial\left(\widetilde{\psi}_1 + \widetilde{\psi}_2\right) = \widetilde{\phi}_1 d\bar z_1 + \widetilde{\phi}_2 d\bar z_2,$$

and

$$(\phi_*)^* = \phi.$$

In the proposition below we identify the spaces of residual cohomologies $H^{(0,1)}_R(\mathcal{C})$ of curve $\mathcal{C}$ and the cohomologies $H^1(\mathcal{C}, \mathcal{O})$ of the structural sheaf $\mathcal{O}$ on $\mathcal{C}$.

**Proposition 3.1.** *(compare with* [Gr, Ha2]*) (Isomorphism of cohomologies) Let*

$$\mathcal{C} = \left\{z \in \mathbb{CP}^2 : P(z) = 0\right\}$$

*be a curve in $\mathbb{CP}^2$ with nodal points as the only singularities. Then there exists an isomorphism*

$$\imath : H^{(0,1)}_R(\mathcal{C}) \to H^1(\mathcal{C}, \mathcal{O}). \tag{3.3}$$

*Proof.* To define $\imath$ we consider an arbitrary $\phi \in Z^1_R(\mathcal{C}, \mathcal{O})$ represented locally on a cover $\{U_i\}_{i=1}^N$ by residual currents

$$\left\{\frac{\Phi^{(0,1)}_i}{F^{(i)}}\right\}_{i=1}^N$$

satisfying $\bar\partial\Phi_i = F^{(i)}\Omega^{(0,2)}_i$. Then, using existence of $\Psi^{(0,1)}_i \in \mathcal{E}^{(0,1)}(U_i)$ such that

$$\bar\partial\Psi^{(0,1)}_i = \Omega^{(0,2)}_i$$

we obtain

$$\bar\partial\left(\Phi_i - F^{(i)}\Psi_i\right) = 0 \ \text{ in } U_i,$$

and therefore

$$\Phi_i = \bar\partial\Theta_i + F^{(i)}\Psi_i$$

for some $\Theta_i \in \mathcal{E}^{(0,0)}(U_i)$.

From the last equality we obtain that

$$\bar{\partial}\left(\Theta_i - \Theta_j\right) = F_i \Omega_{ij} \text{ on } U_{ij} = U_i \cap U_j,$$

and therefore, by defining

$$\imath\{\Phi_i\} = \{\Theta_i - \Theta_j\} \tag{3.4}$$

we obtain a cocycle in $Z^1(\mathcal{C}, \mathcal{O})$.

To construct the inverse to $\imath$ we take $\{\Theta_{ij}\}$ - a cocycle in $Z^1(\mathcal{C}, \mathcal{O})$. We consider a partition of unity $\{\vartheta_i\}_{i=1}^N$ subordinate to some open cover $\{U_i\}_{i=1}^N$ of an open neighborhood $U \supset V$. Without loss of generality we may assume that $\{\Theta_{ij}\}$ are restrictions of holomorphic functions on $U_{ij}$. We define the cochain

$$\Theta_i = \sum_{k \neq i,\ U_k \cap U_i \neq \emptyset} \vartheta_k \Theta_{ik},$$

and notice that on $U_{ij}$ we have the equality

$$\begin{aligned}
\Theta_i - \Theta_j = \sum_{k \neq i,\ U_k \cap U_i \neq \emptyset} \vartheta_k \Theta_{ik} - \sum_{k \neq j,\ U_k \cap U_j \neq \emptyset} \vartheta_k \Theta_{jk} \\
= \sum_{k \neq i,j\ U_k \cap U_i \neq \emptyset} \vartheta_k \left(\Theta_{ik} - \Theta_{jk}\right) + \vartheta_j \Theta_{ij} - \vartheta_i \Theta_{ji} \\
= \sum_{k \neq i,j\ U_k \cap U_i \neq \emptyset} \vartheta_k \Theta_{ij} + \vartheta_j \Theta_{ij} + \vartheta_i \Theta_{ij} + F_i \Omega_{ij} = \Theta_{ij} + F_i \Omega_{ij}
\end{aligned}$$

with some functions $\Omega_{ij} \in \mathcal{E}^{(0,0)}(U_{ij})$, and its corollary

$$\bar{\partial}\left(\Theta_i - \Theta_j\right) = F_i \bar{\partial} \Omega_{ij}. \tag{3.5}$$

Therefore, the collection

$$\left\{ \Phi_i = \bar{\partial} \Theta_i = \sum_{k \neq i,\ U_k \cap U_i \neq \emptyset} \Theta_{ik} \bar{\partial} \vartheta_k \right\}_{i=1}^N \in Z_R^{(0,1)}(\mathcal{C}) \tag{3.6}$$

defines a $\bar{\partial}$-closed residual current in a neighborhood of $V$.

If we apply the map $\imath$ from (3.4) to the current in (3.6), then using equality (3.5) we obtain

$$\imath\{\Phi_i\} = \{\Theta_i - \Theta_j\} = \{\Theta_{ij}\} \in Z^1(\mathcal{C}, \mathcal{O}).$$

□

Let now $\{p^{(i)}\}_{i=1}^r$ be the nodal points in $\mathcal{C}$, and let points $p_1^{(i)}, p_2^{(i)} \in \mathcal{X}$ be such that $\varrho(p_1^{(i)}) = \varrho(p_2^{(i)}) = p^{(i)} \in \mathcal{C}$. We consider a collection of paths $\{\gamma_i\}_{i=1}^r$ in $\mathcal{X}$ such that $\gamma_i(0) = p_1^{(i)}$ and $\gamma_i(1) = p_2^{(i)}$ and functions $\{f_i\}_{i=1}^r \in \mathcal{E}(\mathcal{X})$ with supports in some neighborhoods $U_i \supset \gamma_i$ such that

$$\begin{cases} f_i(p_1^{(i)}) = 0,\ f_i(p_2^{(i)}) = 1, \\ \partial f_i = 0 \text{ in } V_i \Subset U_i. \end{cases} \tag{3.7}$$

Then for a path $\gamma_i \subset \mathcal{X}$ we have

$$\int_{\gamma_i} \bar{\partial} f_i = f_i(p_2^{(i)}) - f_i(p_1^{(i)}) = 1.$$

For an arbitrary form $\phi \in \mathcal{E}^{(0,1)}(\mathcal{X})$ and the corresponding residual current $\phi_*$ on $\mathcal{C}$ we consider the decomposition on $\mathcal{C}$ that follows from Theorem 1:

$$\phi_* = \bar\partial I[\phi_*] + L[\phi_*],$$

and the lift of this decomposition on $\mathcal{X}$:

$$\phi = (\phi_*)^* = \bar\partial (I[\phi_*])^* + (L[\phi_*])^*. \tag{3.8}$$

From Proposition 4.4 in [HP4] it follows that if $\phi \in \mathcal{E}^{(0,1)}(\mathcal{X})$, then $I[\phi_*] \in \mathcal{E}(\mathcal{C})$, and therefore

$$(I[\phi_*])^* \in \mathcal{E}(\mathcal{X}).$$

We consider the scalar product on $\mathcal{E}^{(0,1)}\left(\mathcal{X}\right)$

$$\langle \phi, \psi \rangle = \int_{\mathcal{X}} \phi \wedge \overline{\psi} \tag{3.9}$$

and assume without loss of generality that the collection of forms $\left\{(L[(\bar\partial f_i)_*])^*\right\}_{i=1}^r$ is orthonormalized with respect to scalar product in (3.9). Then for $\phi \in \mathcal{E}^{(0,1)}\left(\mathcal{X}\right)$ we define for $i = 1, \dots, r$,

$$a_i[\phi] = \int_{\mathcal{X}} \phi \wedge \overline{(L[(\bar\partial f_i)_*])^*}, \tag{3.10}$$

and consider operators $\mathcal{L} : \mathcal{E}^{(0,1)}(\mathcal{X}) \to \mathcal{E}^{(0,1)}(\mathcal{X})$ and $\mathcal{I} : \mathcal{E}^{(0,1)}(\mathcal{X}) \to \mathcal{E}(\mathcal{X})$, defined as

$$\mathcal{L}[\phi] = (L[\phi_*])^* - \sum_{i=1}^r a_i[\phi](L[(\bar\partial f_i)_*])^*, \tag{3.11}$$

and

$$\mathcal{I}[\phi] = \left(I\left[\phi_* - \sum_{i=1}^r a_i[\phi](\bar\partial f_i)_*\right]\right)^* + \sum_{i=1}^r a_i[\phi] f_i. \tag{3.12}$$

**Proposition 3.2.** *(Hodge-type decomposition on a nonsingular curve) Let $\mathcal{X}$ be a nonsingular curve, and let $\mathcal{X} \overset{\varrho}{\to} \mathbb{CP}^2$ be an immersion of $\mathcal{X}$ into $\mathbb{CP}^2$ with r nodal points. Let $\{f_i\}_{i=1}^r$ satisfy* (3.7) *and let operators $\mathcal{L}$ and $\mathcal{I}$ be defined respectively in* (3.11) *and* (3.12).

*Then for the space of* $(0,1)$*-forms* $\mathcal{E}^{(0,1)}(\mathcal{X})$

(i) *the following decomposition holds*

$$\phi = \bar\partial \mathcal{I}[\phi] + \mathcal{L}[\phi], \tag{3.13}$$

(ii) *a* $(0,1)$*-form* $\phi \in \mathcal{E}^{(0,1)}(\mathcal{X})$ *is* $\bar\partial$*-exact, iff* $\mathcal{L}[\phi] = 0$.

*Proof.* To prove equality (3.13) we use equality (3.8) to obtain for an arbitrary $\bar\partial$-closed form $\phi \in \mathcal{E}^{(0,1)}(\mathcal{X})$ the equality

$$\phi - \sum_{i=1}^r a_i[\phi] \bar\partial f_i = \bar\partial \left(I\left[\phi_* - \sum_{i=1}^r a_i[\phi](\bar\partial f_i)_*\right]\right)^* + \left(L\left[\phi_* - \sum_{i=1}^r a_i[\phi](\bar\partial f_i)_*\right]\right)^*,$$

which we can rewrite as

$$\phi = \bar\partial \left[\left(I\left[\phi_* - \sum_{i=1}^r a_i[\phi](\bar\partial f_i)_*\right]\right)^* + \sum_{i=1}^r a_i[\phi] f_i\right] + \mathcal{L}[\phi] = \bar\partial \mathcal{I}[\phi] + \mathcal{L}[\phi].$$

According to Theorem 1 and Proposition 3.1 the image of $L^*$ is a subspace of the space of $\bar\partial$-closed forms - $\mathcal{E}^{(0,1)}(\mathcal{X})$ of dimension $\dim H^1_R(\mathcal{C}, \mathcal{O}) = p_a(\mathcal{C})$ - the arithmetic genus of $\mathcal{C}$. We notice that for any collection $\{c_i\}_{i=1}^r$ there is no $g \in \mathcal{E}(\mathcal{C})$ such that

$$\bar\partial g^* = \sum_{i=1}^r c_i \bar\partial f_i,$$

because otherwise we would have

$$g^* - \sum_{i=1}^{r} c_i f_i = \text{const},$$

which contradicts $g^*$ taking the same values at $p_1^{(i)}$ and $p_2^{(i)}$ for all $i = 1, \dots, r$. Therefore,

$$\dim \operatorname{Ker}\mathcal{L} = \dim \operatorname{Span}\left\{(L[(\bar\partial f_i)_*])^*\right\} = r,$$

and

$$\dim \operatorname{Im}\{\mathcal{L}\} = \dim \operatorname{Im}\{L^*\} - r = p_a(\mathcal{C}) - r.$$

To prove item (ii) of Proposition 3.2 we consider a $\bar\partial$-exact form $\phi = \bar\partial g$ for $g \in \mathcal{E}(\mathcal{X})$ and assume that

$$\left(L\left[(\bar\partial g)_* - \sum_{i=1}^{r} a_i[\bar\partial g](\bar\partial f_i)_*\right]\right)^* \neq 0.$$

Then from the inequality above it follows that

$$\dim\left\{\mathcal{L}\left\{\bar\partial g, g \in \mathcal{E}(\mathcal{X})\right\}\right\} > r,$$

which, according to Proposition 3.1, contradicts the statement from IV.1 in [Ha2], that

$$\dim H^{(0,1)}(\mathcal{X}) = p_a(\mathcal{C}) - r.$$

□

Using Theorem 1 and Proposition 3.2 we prove the following version of the Hodge Theorem for smooth algebraic curves, which gives an explicit formula for solution $R_1[\phi]$ of equation

$$\bar\partial R_1[\phi] = \phi - H_1[\phi].$$

**Theorem 2.** *(Explicit formulas in the Hodge decomposition) Let $\mathcal{X}$ be a smooth algebraic curve, let $\mathcal{L}$ and $\mathcal{I}$ be operators from* (3.11) *and* (3.12) *respectively, and let $\{\omega_j\}_{j=1}^{g}$ be an orthonormal basis of holomorphic* $(1,0)$*-forms on $\mathcal{X}$, i.e.*

$$\int_{\mathcal{X}} \omega_j \wedge \bar\omega_k = \delta_{jk},\ j,k = 1, \dots, g.$$

*Then Hodge operators $H_1$ and $R_1$ in decomposition* (1.1) *admit the following representations*

$$\begin{aligned} H_1[\phi] &= \sum_{j=1}^{g}\left(\int_{\mathcal{C}} \phi \wedge \omega_j\right)\bar\omega_j, \\ R_1[\phi] &= \mathcal{I}\big[\phi + (\mathcal{L} - H_1)[\phi]\big] + \textit{const}. \end{aligned} \tag{3.14}$$

*Proof.* Combining decompositions (1.1) and (2.9) we obtain equality

$$\bar\partial\big(R_1[\phi] - \mathcal{I}[\phi]\big) = (\mathcal{L} - H_1)[\phi]. \tag{3.15}$$

Then, applying Theorem1 to the $\bar\partial$-exact form $\psi = (\mathcal{L} - H_1)[\phi]$ we obtain equalities

$$\psi = \bar\partial\mathcal{I}[\psi]$$

and

$$(\mathcal{L} - H_1)[\phi] = \bar\partial\big(\mathcal{I}(\mathcal{L} - H_1)[\phi]\big). \tag{3.16}$$

Finally, from equalities (3.15) and (3.16) we obtain equality

$$\bar\partial\big(R_1[\phi] - \mathcal{I}[\phi] - \mathcal{I}(\mathcal{L} - H_1)[\phi]\big) = 0,$$

which implies the second equality in (3.14). □

## 4. EXPLICIT SOLUTION OF $\bar\partial$-EQUATION ON AFFINE CURVES.

In this section we prove solvability of the $\bar\partial$-equation on affine smooth algebraic curves. Let $\mathcal{X}$ be a nonsingular algebraic curve, and let $\varrho : \mathcal{X} \to \mathbb{CP}^2$ be an immersion of $\mathcal{X}$ as in (3.1) such that

$$\mathcal{C} = \varrho(\mathcal{X}) = \left\{z \in \mathbb{CP}^2 : P(z) = 0\right\} \tag{4.1}$$

is a plane curve of $\deg P = d$ with at most nodes as singularities. Without loss of generality we may assume that the intersection of $\mathcal{C}$ with the line at infinity

$$\mathcal{C} \cap \left\{(z_0, z_1, z_2) \in \mathbb{CP}^2 : z_0 = 0\right\} = \left\{z^{(1)}, \dots, z^{(r)}\right\} \tag{4.2}$$

consists of $r$ points with multiplicities $m_1, \dots, m_r$ such that $\sum_{i=1}^{r} m_i = d$. We denote

$$\mathring{\mathcal{C}} = \mathcal{C} \setminus \left\{z^{(1)}, \dots, z^{(r)}\right\},$$

and

$$\mathring{\mathcal{X}} = \mathcal{X} \setminus \left\{\varrho^{-1}(z^{(1)}), \dots, \varrho^{-1}(z^{(r)})\right\}.$$

**Proposition 4.1.** *(Hodge-Kodaira Vanishing Theorem for affine curves) Let $\mathcal{X}$ and $\mathcal{C} \subset \mathbb{CP}^2$ be curves as in* (4.1)*, with $\mathcal{C}$ of degree $d$ satisfying condition* (4.2)*. Let $\phi^{(0,1)} \in \mathcal{E}(\mathring{\mathcal{X}})$ be a form on $\mathring{\mathcal{X}}$, and $\phi_*$ be its direct image on $\mathring{\mathcal{C}}$. If for some $\ell$ satisfying $\ell > d - 3$ the form $\zeta_0^{\ell}\phi_*(\zeta)$ admits an extension $\psi(\zeta) \in C^{(0,1)}(\mathcal{C})$, then there exists a function $g \in \mathcal{E}(\mathring{\mathcal{X}})$ such that*

$$\begin{cases} \bar\partial g = \phi, \\ |g(\zeta)| \leq C|\zeta_0|^{-\ell}. \end{cases} \tag{4.3}$$

*Proof.* Let $z^{(j)} \in \mathcal{C} \cap \left\{\mathbb{CP}^2 : z_0 = 0\right\}$ be one of the points of $\mathcal{C}$ at infinity. We consider a neighborhood $U^{(j)}$ of $z^{(j)}$ in $\mathbb{C}^2$ with coordinates $\zeta_1, \zeta_2$ such that

$$\mathcal{C} \cap U^{(j)} = \left\{\zeta \in U^{(j)} : \zeta_2 = 0\right\}.$$

Using Cauchy-Green formula we solve the $\bar\partial$-equation on $\mathcal{C} \cap U^{(j)}$ and obtain a function

$$g^{(j)}(\zeta_1) \in C(\mathcal{C}) \cap C^{\infty}(\mathring{\mathcal{C}})$$

with compact support in $\mathcal{C} \cap U^{(j)}$ satisfying equality

$$\bar\partial g^{(j)} = \psi\big|_{\mathcal{C} \cap V^{(j)}},$$

where $V^{(j)} \Subset U^{(j)}$.

Then the form

$$\tilde\psi = \psi - \sum_{j=1}^{d} \bar\partial g^{(j)}$$

defines a $\bar\partial$-closed residual current on $\mathcal{C}$ of degree $\ell > d - 3$, and therefore, using Corollary 2.4 we obtain the existence of a function $\tilde g = I_1[\zeta_0^{\ell} \cdot \tilde\psi(\zeta)] \in C^{(0,0)}(\mathring{\mathcal{C}})$ satisfying the equation

$$\bar\partial \tilde g = \tilde\psi.$$

Therefore function $g_* = \tilde g + \sum_{j=1}^{d} g^{(j)}$ satisfies the equation

$$\bar\partial g_*(\zeta) = \psi(\zeta) = \zeta_0^{\ell}\phi_*(\zeta),$$

and the function $g(\zeta) = (\zeta_0^{-\ell} g_*)^*$ satisfies conditions (4.3). □

Combining results of Theorem 2 and Proposition 4.1 we obtain the following proposition.

**Proposition 4.2.** *Let* $\mathcal{X}$ *and* $\mathcal{C} \subset \mathbb{CP}^2$ *be curves as in* (4.1)*, with* $\mathcal{C}$ *satisfying condition* (4.2)*. Let* $\phi^{(0,1)} \in \mathcal{E}(\mathring{\mathcal{X}})$ *be a form on* $\mathring{\mathcal{X}}$*, such that its direct image* $\phi_*$ *has a compact support in* $\mathring{\mathcal{C}}$*. Then for arbitrary* $\ell > d-3$ *the function*

$$g(\zeta) = \left(\zeta_0^{-\ell} \cdot I_1[\zeta_0^{\ell}\phi_* - H_1[\zeta_0^{\ell}\phi_*]]\right)^*$$

*satisfies equation*

$$\bar{\partial} g = \phi.$$

## REFERENCES

[AS] M. Andersson, H. Samuelsson, A Dolbeault-Grothendieck lemma on complex spaces via Koppelman formulas, Inv. Math. (2012) 190, 261-297.

[Bn] B. Berndtsson, Integral formulas on projective space and the Radon transform of Gindikin-Henkin-Polyakov, Publ. Math. Universitat Autonoma de Barcelona, 32(1) (1988), 7-41.

[BDIP] J. Bertin, J-P. Demailly, L. Illusie, C. Peters, Introduction to Hodge Theory, SMF/AMS TEXTS and MONOGRAPHS, V. 8, 2002.

[C] A.P. Calderon, On an inverse boundary problem, In Seminar on Numerical Analysis and Its Applications to Continuum Physics, Soc. Brasiliera de Matematica, (1980), 61-73.

[CH] N.R. Coleff, M.E. Herrera, Les Courants Résiduels Associés à une Forme Méromorphe, Lecture Notes in Mathematics 633, Springer Verlag, New York, 1978.

[Dl] P. Deligne, Théorie de Hodge I, II, III, Actes du Congrès International des Mathématiciens (Nice, 1970), Tome 1, 425-430, Gauthier-Villars, Paris, 1971, Inst. Hautes Études Sci. Publ. Math. No. 40 (1971), 5-57, No. 44 (1974), 5-77.

[Do] P. Dolbeault, Theorie des résidus, Seminar P. Lelong, 1970/71, Exp. 09/08/1970.

[FG] J. E. Fornæss, E. A. Gavosto, The Cauchy-Riemann equation on singular spaces, Duke Math. J. 93(3) (1998), 453477.

[Ga] C.F. Gauss, General investigations of curved surfaces of 1827 and 1825, C. S. Robinson & Co., Princeton University Press, 1902.

[Ge] I.M. Gelfand, Some problems of Functional Analysis and Algebra, in Proc. Int. Congr. Math. (Amsterdam 1954), 253-276.

[GH] Ph. Griffiths, J. Harris, Principles of algebraic geometry, John Wiley and Sons, New York, 1978.

[Gr] A. Grothendieck, The cohomology theory of abstract algebraic varieties, Proc. Internat. Congress Math. (Edinburgh, 1958), 103-118, 1960, Cambridge Univ. Press, New York.

[Ha1] R. Hartshorne, Residues and duality, Lecture Notes in Mathematics 20, Springer Verlag, 1966.

[Ha2] R. Hartshorne, Algebraic Geometry, Springer-Verlag, New York, 1977.

[HL] M. Herrera, D. Lieberman, Residues and principal values on complex spaces, Math. Ann. 194, (1971) 259-294.

[HN] G.M. Henkin, R. G. Novikov, On the reconstruction of conductivity of a bordered two-dimensional surface in $\mathbb{R}^3$ from electrical current measurements, on its boundary, J. Geom. Anal. 21 (2011), no. 3, 543-587.

[HP1] G.M. Henkin, P.L. Polyakov, Homotopy formulas for the $\bar{\partial}$-operator on $\mathbb{CP}^n$ and the Radon-Penrose transform, Izv. Akad. Nauk SSSR Ser. Mat. 50:3 (1986), 566-597.

[HP2] G.M. Henkin, P.L. Polyakov, The Grothendieck-Dolbeault lemma for complete intersections, C. R. Acad. Sci., Ser I, Math. 308 (1989), 405-409.

[HP3] G.M. Henkin, P.L. Polyakov, Residual $\bar{\partial}$-cohomology and the complex Radon transform on subvarieties of $\mathbb{CP}^n$, Math. Ann. 354:2 (2012), 497-527.

[HP4] G.M. Henkin, P.L. Polyakov, Explicit Hodge-type decomposition on projective complete intersections, The Journal of Geometric Analysis, 26(1), 2016, 672-713, DOI 10.1007/s12220-015-9643-1.

[Ho] W.V.D. Hodge, The theory and applications of harmonic Integrals, 2-nd ed., Cambridge University Press, 1952.

[Kl] F.Klein, On Riemann's theory of algebraic functions and their integrals, Macmillan and Bowes, Cambridge, 1893.

[Kd1] K. Kodaira, Harmonic fields in Riemannian manifolds, Ann. of Math., 50 (1949), 587-665.

[Kd2] K. Kodaira, On Käler varieties of restricted type, Ann. of Math., 60 (1954), 28-48.

[Kp] W. Koppelman, The Cauchy integral for differential forms, Bull. AMS 73 (1967), 554-556.

[L] J. Leray, Le calcul differéntiel et intégral sur une variété analytique complexe, Bull. Soc. Math. France 87 (1959), 81-180.

[Li] I. Lieb, Die Cauchy-Riemannschen Differentialgleichungen auf streng pseudokonvexen Gebieten, Math. Ann. 190 (1970), 6-44.

[O] N. Øvrelid, Integral representation formulas and $L_p$-estimates for the $\bar{\partial}$-equation, Math. Scand. 29 (1971), 137-160.

[P] H. Poincaré , Sur les résidus des intégrales doubles, Acta Math. 9, (1887), 321-380.

[Pa] M. Passare, Residues, currents, and their relation to ideals of holomorphic functions, Math. Scand. 62 (1988), 75-152.

[Po] P. L. Polyakov, Cauchy-Weil formula for differential forms, Mat. Sb. (N.S.) 85:3 (1971), 383-398.

[RR] J.P. Ramis, G. Ruget, Complexe dualisant et théorèmes de dualité en géométrie analytique complexe, Publ. Math. de l'I.H.E.S., 38 (1970), 77-91.

[Sa] H. Samuelson Kalm, Holomorphic forms, the $\bar{\partial}$-equation, and duality on a reduced complex space, Preprint, 2015.

[Sp] G. Springer, Introduction to Riemann surfaces, Addison-Wesley, Reading, Ma,1957.

[T] Y.L. Tong, Integral representation formulae and Grothendieck residue symbol, Amer. J. Math., 95:4, (1973), 904-917.

[Wi] A. Weil, L'intégrale de Cauchy et les fonctions de plusieurs variables, Math. Ann. 111 (1935), no. 1, 178-182.

[Wy1] H. Weyl, The method of orthogonal projection in potential theory, Duke Math. J. 7, (1940), 411-444.

[Wy2] H. Weyl, On Hodge's theory of harmonic integrals, Ann. of Math. (2) 44, (1943), 1-6.

INSTITUT DE MATHEMATIQUES, UNIVERSITE PIERRE ET MARIE CURIE, 75252 BC247 PARIS, CEDEX 05, FRANCE, AND CEMI ACAD. SC., 117418, MOSCOW, RUSSIA

*E-mail address*: `guennadi.henkin@imj-prg.fr`

UNIVERSITY OF WYOMING, DEPARTMENT OF MATHEMATICS, 1000 E UNIVERSITY AVE, LARAMIE, WY 82071

*E-mail address*: `polyakov@uwyo.edu`